

RONA Z. SILKISS

M.D., FACS

OCULOFACIAL PLASTIC, RECONSTRUCTIVE AND ORBITAL SURGERY

ADMINISTRATIVE OFFICE  
400 29th Street, Suite 315  
Oakland, CA 94609  
Phone 510.763.0881  
Fax 510.763.0907

## Latisse Ophthalmic Solution Consent

Please read the following information carefully. Although minimal, there are possible side effects and contraindications to the use of Latisse.

You should not use Latisse if you are allergic to any of its ingredients. You should inform the doctor if you:

- \* Have been previously diagnosed with eye pressure problems
- \* Are currently taking medication for eye pressure problems
- \* Have risk factors for Glaucoma

The most common side effects of Latisse are itching or redness. These were reported by 4% of users in the clinical trial. Latisse use may cause other less common side effects such as, darkening of the eye lid, skin, eye irritation, dryness or redness of the eyelids or increased brown pigmentation of the colored part of the eye (iris). The pigmentation of the iris may be permanent.

If you develop a new eye condition (ie, eye trauma or infection) , experience a sudden decrease in vision, have eye surgery, or develop any eye or eyelid reactions , you should immediately discontinue use of Latisse and talk to your doctor.

My signature below constitutes my full understanding of the risks and side effects of the prescription solution, Latisse.

Signature \_\_\_\_\_

Date \_\_\_\_\_