

Silkiss Eye Surgery
400 - 29th Street, Suite 315, Oakland, CA 94609
Phone 510.763.0881 / Fax 510.763.0907

Please do not ingest aspirin, or nonsteroidal anti-inflammatory medication for two weeks prior to surgery. If you are on Coumadin, Plavix, Aggrenox, or other blood-thinning medication, examples below, please consult with your physician to determine when this medication should be discontinued prior to surgery- it may be less than two weeks. You may restart blood-thinners, aspirin, and nonsteroidal anti-inflammatory medications **the day after** surgery. You should continue to take your other medications on your usual schedule.

Advil	Empirin	Savaysa (Edoxaban)
Aggrenox	Empirazil	Sleeping Aid
Aleve	Equagesic	St. Joseph's Cold Tabs
Alka Seltzer	Excedrin	Sine-Off Cold Tablets
Anacin	Fiogesic	Stanback Analgesic Powders
Anaprox	Fiorinal	Supac
APC	Fish Oil	Synalgos
Arthritis Pain Formula	Flaxseed Oil	Ticlid
Arthritis Strength Bufferin	4-Way Cold Tablets	Triaminicin Tablets
Ascriptin	Goddy's Headache Powder	Vanquish
Aspirin	Ibuprofen	Vitro-Med Tablets
Aspirin Suppositories	Indocin	Vioxx
Bayer Children's Cold Tabs	Midol	Voltaren
Brilinta	Motrin	Warfarin
Buff-A-Comp	Naprosyn	Xarelto (Rivaroxaban) 2-3 days
Bufferin	Norgesic	Zorprin
Cama Inlay Tabs	Omega 3 Fatty Acids	
Celebrex	Pabirin	<u>HERBS</u>
Clopidogrel	Pacaps	Certain Chinese Herbs
Congesperin	Panalgesic	Garlic Pills
Cordicin Products	Pedrazil	Ginseng Pills
Coumadin 5 days	Percodan	Ginkgo Pills
Darvon with ASA	Persistin	Ginger Pills
Dristan	Plavix 5-7 days	
Ecotrin	Pradaxa (Dabigatran)	Tylenol is ok to take prior to surgery.
Edoxaban	Quiet World Analgesic	
Eliquis (Apixaban)	Robaxisal	

New diabetes and weight loss medications may also need to be stopped prior to surgery

GLP Agonists need to be stopped prior to surgery.

If you take a weekly dose, stop one week prior to surgery.

If you take a daily dose, stop one day prior to surgery.

Dulaglutide (Trulicity) (injection weekly)
Exenatide extended release (Bydureon) (injection weekly)
Exenatide (Byetta) (injection twice daily)
Semaglutide (Ozempic) (injection weekly)
Semaglutide (Rybelsus) (taken by mouth once daily)
Semaglutide (Wegovy) (injection weekly)
Liraglutide (Victoza, Saxenda) (injection once daily)
Lixisenatide (Adlyxin) (injection once daily)
Tirzepadite (Mounjaro) (injection weekly)

Oral SGLT2 Inhibitors need to be stopped prior to surgery with general anesthesia.

Bexagliflozin (Brenzavvy) STOP 3 days prior to surgery.
Canagliflozin (Invokana) STOP 3 days prior to surgery.
Dapagliflozin (Farxiga) STOP 3 days prior to surgery.
Empagliflozin (Jardiance) STOP 3 days prior to surgery.
Ertugliflozin (Steglatro) STOP 4 days prior to surgery.